

# INTERSECTION COHOMOLOGY, MONODROMY, AND THE MILNOR FIBER

DAVID B. MASSEY

ABSTRACT. We say that a complex analytic space,  $X$ , is an intersection cohomology manifold if and only if the shifted constant sheaf on  $X$  is isomorphic to intersection cohomology; this is quickly seen to be equivalent to  $X$  being a homology manifold. Given an analytic function  $f$  on an intersection cohomology manifold, we describe a simple relation between  $V(f)$  being an intersection cohomology manifold and the vanishing cycle Milnor monodromy of  $f$ . We then describe how the Sebastiani-Thom isomorphism allows us to easily produce intersection cohomology manifolds with arbitrary singular sets. Finally, as an easy application, we obtain restrictions on the cohomology of the Milnor fiber of a hypersurface with a special type of one-dimensional critical locus.

## §0. Introduction

Suppose that  $X$  is a connected open neighborhood of the origin in  $\mathbb{C}^n$ , where  $n \geq 2$ , and suppose that  $f : X \rightarrow \mathbb{C}$  is (complex) analytic. Let  $\mathbf{p} \in V(f)$  and let  $F_{f,\mathbf{p}}$  denote the Milnor fiber of  $f$  at  $\mathbf{p}$ .

In Theorem 8.5 of Milnor's classic "Singular Points of Complex Hypersurfaces" [Mi], Milnor proved that, if  $n \neq 3$  and  $f$  has an isolated critical point at  $\mathbf{p}$ , then the real link of  $V(f)$  at  $\mathbf{p}$  is a topological sphere if and only if the identity minus the (Milnor) monodromy, acting on  $H^{n-1}(F_{f,\mathbf{p}}; \mathbb{Z})$ , is invertible over  $\mathbb{Z}$ .

Milnor's proof can easily be used in the case where  $f$  may have non-isolated critical points; in this setting, Milnor's proof shows that  $V(f)$  is an integral cohomology manifold (or integral homology manifold) if and only if, for all  $\mathbf{p} \in V(f)$ , the identity minus the monodromy,  $\text{id} - \tilde{T}_{f,\mathbf{p}}$ , acting on the reduced cohomology of the Milnor fiber of  $f$  at  $\mathbf{p}$ ,  $\tilde{H}^*(F_{f,\mathbf{p}}; k)$ , induces isomorphisms (in all degrees). This was demonstrated in 1977 by Randell [R]. Our version of this result appears in Theorem 2.2.

Let us work now over a base ring  $k$  which is actually a field (as opposed to working over  $\mathbb{Z}$ ). The categorical implication of the previous paragraph is that  $V(f)$  is a  $k$ -cohomology manifold if and only if the morphism  $\text{id} - \tilde{T}_f$ , acting on the vanishing cycles  $\phi_f[-1]k_x^\bullet[n]$ , is an isomorphism in the derived category  $D_c^b(V(f))$  and, hence, an isomorphism in the full subcategory  $\text{Perv}(V(f))$  of perverse sheaves on  $V(f)$ . This leads one to ask: is saying that an analytic space  $Y$  is a cohomology or homology manifold equivalent to some nice statement in the category  $\text{Perv}(Y)$ ?

As Borho and MacPherson showed in [B-M] and as we recall in Theorem 1.1, the answer to the above question, is "yes". In fact, the axiomatic characterization of intersection cohomology in terms of the support and co-support conditions implies immediately that an  $m$ -dimensional complex analytic space  $Y$  is a  $k$ -cohomology manifold if and only if the shifted constant sheaf  $k_Y^\bullet[m]$  is isomorphic to the (middle perversity) intersection cohomology complex  $IC_Y^\bullet$ . Here, by intersection cohomology, we mean that we are using the indexing which places possibly non-zero cohomology only in non-negative degrees, i.e., if  $Y$  were a purely  $m$ -dimensional **manifold**, then  $IC_Y^\bullet$  would be isomorphic to the shifted constant sheaf  $k_Y^\bullet[m]$ .

---

2000 *Mathematics Subject Classification.* 32B15, 32C35, 32C18, 32B10.

*Key words and phrases.* intersection cohomology, monodromy, Milnor fiber, vanishing cycles.

What is the importance of  $k_Y^\bullet[m]$  being isomorphic to  $IC_Y^\bullet$ ? It is of great categorical importance, for if  $Y$  is irreducible, then  $IC_Y^\bullet$  is a simple object in the Abelian, locally Artinian category  $Perv(Y)$ . Thus, if  $Y$  is irreducible,  $Y$  is a  $k$ -cohomology manifold if and only if  $k_Y^\bullet[m]$  is a simple object in  $Perv(Y)$ . As it is this categorical property of cohomology manifolds that we make use of throughout our arguments, we prefer to refer to such spaces  $Y$  not as  $k$ -cohomology manifolds, but, rather, as  *$k$ -intersection cohomology manifolds* (or  *$k$ - $IC^\bullet$  manifolds*).

In Theorem 2.2, we demonstrate the usefulness of the categorical approach. We consider  $f : X \rightarrow \mathbb{C}$ , but no longer require  $X$  to be smooth – instead, we allow  $X$  itself to be a  $k$ - $IC^\bullet$  manifold. In this generalized setting, we relate  $V(f)$  being a  $k$ - $IC^\bullet$  manifold to  $\text{id} - \tilde{T}_f$  being an isomorphism in  $Perv(V(f))$ . The proof that we give is very short (though significant knowledge of the derived category is required); moreover, the hypotheses of the theorem are as weak as possible and are natural from the categorical point-of-view. Much of Theorem 2.2 could be proved along the lines of Milnor’s argument, but the argument would be longer, special cases would be problematic, and the optimal hypotheses would be difficult to arrive at.

Let us consider a simple, but not trivial, example:  $k = \mathbb{C}$ ,  $X = \mathbb{C}^2$ , and  $f : X \rightarrow \mathbb{C}$  is given by  $f(xy) = xy$ . Let  $m$  denote the inclusion of the origin,  $\mathbf{0}$ , into  $V(f)$ . Let  $j^x$  (respectively,  $j^y$ ) denote the inclusion of  $V(y)$  (respectively,  $V(x)$ ) into  $V(f)$ .

It is well-known that the reduced cohomology (with  $k$ -coefficients) of  $F_{f,\mathbf{0}}$  is non-zero only in degree 1,  $\tilde{H}^1(F_{f,\mathbf{0}}) \cong \mathbb{C}$ , and  $\tilde{T}_{f,\mathbf{0}}^1$  is the identity. Hence,  $\phi_f[-1]\mathbb{C}_X^\bullet[2]$  is simply the extension by zero of the constant sheaf at the origin, i.e.,  $\phi_f[-1]\mathbb{C}_X^\bullet[2] \cong m_!\mathbb{C}_\mathbf{0}^\bullet \cong m_*\mathbb{C}_\mathbf{0}^\bullet$ , and  $\text{id} - \tilde{T}_f = 0$ .

This agrees with Theorem 2.2, since

$$IC_{V(xy)}^\bullet \cong j_!^x \mathbb{C}_{V(y)}^\bullet[1] \oplus j_!^y \mathbb{C}_{V(x)}^\bullet[1] \not\cong \mathbb{C}_{V(f)}^\bullet[1].$$

There is another interesting aspect of this example. Consider the nearby cycles  $\psi_f[-1]\mathbb{C}_X^\bullet[2]$  and their monodromy  $T_f$ . Then, the triple  $(\text{id}, T_f, \tilde{T}_f)$  acts on the canonical short exact sequence of perverse sheaves

$$0 \rightarrow C_{V(f)}^\bullet[1] \rightarrow \psi_f[-1]\mathbb{C}_X^\bullet[2] \rightarrow \phi_f[-1]\mathbb{C}_X^\bullet[2] \rightarrow 0.$$

Since  $\tilde{T}_f = \text{id}$ , it follows that  $(\text{id} - T_f)^2 = 0$ . However, it is not difficult to prove that, in a much more general setting, the image of  $\text{id} - T_f$  is isomorphic to  $\phi_f[-1]\mathbb{C}_X^\bullet[2]$ . Thus,  $T_f \neq \text{id}$ . Therefore,  $T_f = \text{id} + N$ , where  $N \neq 0$  and  $N^2 = 0$ .

This non-trivial nilpotent portion of  $T_f$  is related to a deep result of Gabber (see [Mac], §13). Gabber’s result, applied to our current simple case, tells us that the existence of a non-trivial nilpotent portion of  $T_f$ , when  $\tilde{T}_f$  has no such non-trivial nilpotent piece, occurs precisely because  $V(xy)$  is not an intersection cohomology manifold.

In Theorem 2.2, we make the above remark precise; we prove that  $V(f)$  is a  $k$ - $IC^\bullet$  manifold if and only if there is an isomorphism of pairs of complexes and automorphisms

$$(\psi_f[-1]k_X^\bullet[n]; T_f) \cong (k_{V(f)}^\bullet[n-1] \oplus \phi_f[-1]k_X^\bullet[n]; \text{id} \oplus \tilde{T}_f).$$

This result tells us that, while, in general, there may be extra structure present in the nearby cycles and their monodromy, in the case where  $V(f)$  is a  $k$ - $IC^\bullet$  manifold, one may as well simply study the vanishing cycles and their monodromy.

Of course, the reader may suspect that if  $V(f)$  has non-isolated singularities, then  $V(f)$  would almost **never** be a  $k$ - $IC^\bullet$  manifold. In fact, by combining Theorem 2.2 with the Sebastiani-Thom

Theorem ([S-T], [O], [Sa], [N1], [N2], [Mas4], [Sch]), we show in Theorem 3.2 that simple modifications of  $f$  will produce a sequence of functions,  $f_j$ , whose zero loci are  $k$ -IC $^\bullet$  manifolds and whose critical loci are isomorphic to the critical locus of  $f$ . (Here, by “critical locus”, we actually mean the support of the vanishing cycles.) The modifications  $f_j$  are so closely related to  $f$  that questions about the vanishing cycles and monodromy for arbitrary functions  $f$  are reduced to the case where one may assume that  $V(f)$  is an intersection cohomology manifold.

In Section 4, we give a simple “application” of the results of Sections 1 and 2 to line singularities. Suppose that the domain of  $f$  is an open neighborhood of the origin in  $\mathbb{C}^n$  and that  $f$  has a one-dimensional critical locus through the origin. By selecting a linear form  $z_0$  so that  $f|_{V(z_0)}$  has an isolated critical point at  $\mathbf{0}$ , we may define a family of isolated singularities  $f_t := f|_{V(z_0-t)}$ . It is unknown, in the general case, how to effectively calculate the cohomology of the Milnor fiber,  $F_{f,\mathbf{0}}$ , of  $f$  at the origin. (However, many very nice results are known; in particular, the work of Barlet in [B] is very closely related to the results of this paper.)

In Theorem 4.1, we recall that the reduced cohomology  $\tilde{H}^{n-2}(F_{f,\mathbf{0}})$  is a monodromy-invariant subspace of both  $\tilde{H}^{n-2}(F_{f_0,\mathbf{0}})$  and  $\bigoplus_\nu \tilde{H}^{n-2}(F_{f_{t_0},\mathbf{p}_\nu})$ , where the summation is over all irreducible components  $\nu$  of  $\Sigma f$  through the origin and  $\mathbf{p}_\nu$  is any point in  $\mathring{B}_\epsilon \cap \nu \cap V(z_0 - t_0)$  for  $0 < |t_0| \ll \epsilon \ll 1$ . As a corollary of Theorems 2.2 and 4.1, we conclude that if  $V(f_0)$  is a  $k$ -intersection cohomology manifold and  $V(f_{t_0})$  is not, then  $\tilde{H}^{n-2}(F_{f,\mathbf{0}})$  is a **proper** subspace of  $\bigoplus_\nu \tilde{H}^{n-2}(F_{f_{t_0},\mathbf{p}_\nu})$ .

As an example of this last corollary, consider the Whitney umbrella, presented as nodes degenerating to a cusp, given by  $f = z_2^2 - z_1^3 - z_0 z_1^2$ . As  $f_{t_0}$  has a node at the origin, the shifted constant sheaf on  $V(f_{t_0})$  is not intersection cohomology; intersection cohomology would “count the origin twice”. However, the shifted constant sheaf on  $V(f_0)$  is intersection cohomology, since the cusp is a topological manifold. Therefore,  $\tilde{H}^1(F_{f,\mathbf{0}})$  is a proper subspace of  $\bigoplus_\nu \tilde{H}^{n-2}(F_{f_{t_0},\mathbf{p}_\nu}) \cong \tilde{H}^1(F_{f_{t_0},(t_0,\mathbf{0})}) \cong k$ , and our corollary “explains” the well-known fact that  $\tilde{H}^1(F_{f,\mathbf{0}}) = 0$ .

We should remark here that our results are closely related to the work of many people from 15, 20, or even 40 years ago. The results that we prove in Theorems 1.1, 2.2, and 3.2 are very closely related to work of Milnor, Sebastiani and Thom, Goresky and MacPherson, Borho and MacPherson, Randell, Barlet, Kashiwara, Malgrange, Sabbah, and, no doubt, others. By combining old results of these mathematicians, we could arrive at many of our conclusions fairly quickly. Moreover, recent work of Torrelli also yields some of our results. However, the proofs that we give here are extremely short and do, of course, use background material that was developed during the 1980’s and early 1990’s. Moreover, we need the statements of our theorems in the form and generality that we give.

We would like to thank Mark Goresky, Alberto Verjovsky, Tristan Torrelli, and Daniel Barlet for making a number of helpful remarks.

## §1. Intersection Cohomology Manifolds

Fundamental references for many of the tools used throughout this paper are [G-M1], [G-M2], [G-M3], and [K-S].

Throughout this paper, we let  $X$  be an  $n$ -dimensional complex analytic space, and let  $f : X \rightarrow \mathbb{C}$  be a complex analytic function. By intersection cohomology, we shall always mean middle perversity intersection cohomology.

Fix a field  $k$ . Frequently one wishes to take  $k = \mathbb{Q}$  or  $k = \mathbb{C}$ ; however, to detect torsion, one may wish to take the finite fields  $k = \mathbb{Z}/p\mathbb{Z}$ . All cohomology in this paper is calculated with  $k$ -

coefficients. We use field coefficients for two reasons: to make certain duality statements true, and so that the category of perverse complexes of sheaves of  $k$ -modules (vector spaces) on  $X$ ,  $Perv(X)$ , is a locally Artinian category in which the simple objects are extensions by zero of intersection cohomology complexes on irreducible subvarieties with coefficients in an irreducible local system.

Below, it is convenient to adopt the convention that the empty set is a  $(-1)$ -dimensional  $k$ -cohomology sphere. We let  $\mathcal{D}$  denote the Verdier dualizing functor on the derived category  $D_c^b(X)$ .

The following theorem, over the rational numbers, is due to Borho and MacPherson [B-M]; the proof over arbitrary fields remains the same, and all of the equivalences below are either in the statement or the proof of Proposition 1.4 of [B-M]. We give our own proof of the theorem for the sake of self-containment.

**Theorem 1.1** ([B-M]). *The following are equivalent:*

- 1)  $X$  is a  $k$ -intersection cohomology manifold (i.e.,  $IC_x^\bullet \cong k_x^\bullet[n]$ );
- 2)  $\mathcal{D}(k_x^\bullet[n]) \cong k_x^\bullet[n]$ ;
- 3) for all  $x \in X$ , there is an isomorphism of stalk cohomology  $H^*(\mathcal{D}(k_x^\bullet[n]))_x \cong H^*(k_x^\bullet[n])_x$ ;
- 4)  $X$  is a  $k$ -cohomology (or homology) manifold of real dimension  $2n$ , i.e., for all  $x \in X$ , for all  $i$ ,  $H^i(X, X - \{x\}; k) = 0$  unless  $i = 2n$ , and  $H^{2n}(X, X - \{x\}; k) \cong k$ ;
- 5) for all  $x \in X$ , the real link of  $X$  at  $x$  is a  $(2n - 1)$ -dimensional  $k$ -cohomology sphere;
- 6) there exists a Whitney stratification,  $\mathcal{S}$ , of  $X$  such that, for all  $S \in \mathcal{S}$ , if  $d$  is the complex dimension of  $S$ , then the real link of  $S$  is a  $(2(n - d) - 1)$ -dimensional  $k$ -cohomology sphere;
- 7) for all Whitney stratifications,  $\mathcal{S}$ , of  $X$ , for all  $S \in \mathcal{S}$ , if  $d$  is the complex dimension of  $S$ , then the real link of  $S$  is a  $(2(n - d) - 1)$ -dimensional  $k$ -cohomology sphere;

*Proof.* Intersection cohomology is characterized by two conditions: the support and cosupport conditions (see [G-M2]). The support condition is always satisfied by  $k_x^\bullet[n]$  on an  $n$ -dimensional space. In addition, the support and cosupport conditions are dual to each other when using field coefficients. Moreover, the support condition is a condition on stalk cohomology. Hence, 1), 2), and 3) are equivalent.

Now,  $H^i(\mathcal{D}(k_x^\bullet[n]))_x \cong H^{-i+n}(X, X - \{x\}) \cong H_{-i+n}(X, X - \{x\})$ , where the last isomorphism holds since our coefficients are in a field. Thus, 3) and 4) are equivalent, whether we use homology manifolds or cohomology manifolds.

Locally embed  $X$ , at  $x$ , into affine space and let  $B_\epsilon(x)$  denote a small closed ball, of radius  $\epsilon$ , centered at  $x$ , in the ambient space. Then, by excision,

$$H^{-i+n}(X, X - \{x\}) \cong H^{-i+n}(B_\epsilon(x) \cap X, B_\epsilon(x) \cap X - \{x\}) \cong$$

$$\tilde{H}^{-i+n-1}(B_\epsilon(x) \cap X - \{x\}) \cong \tilde{H}^{-i+n-1}(K_{X,x}),$$

where  $K_{X,x}$  denotes the real link of  $X$  at  $x$ . Thus, 4) and 5) are equivalent.

That 6) and 7) are equivalent follows from the equivalence of 1) and 5), and the fact that a complex is the intersection cohomology complex if and only if the shifted restriction of the complex to normal slices to Whitney strata yields intersection cohomology.  $\square$

*Remark 1.2.* Mark Goresky has pointed out that we do not need complex analytic spaces or middle perversity intersection cohomology for some of the implications in Theorem 1.1; if  $X$  is a reasonable space (say, homeomorphic to a simplicial complex) and the shifted constant sheaf on  $X$  is isomorphic to the intersection cohomology complex, with **any** perversity, then  $X$  is a  $k$ -homology manifold.

In [T], in the case where  $X$  is a local complete intersection, Tristan Torrelli gives another characterization for  $X$  being a  $\mathbb{C}$ - $IC^\bullet$  manifold; this characterization is in terms of the Bernstein functional equation. See also Remark 2.4 for the hypersurface case.

**Corollary 1.3.** *Suppose that  $X$  is an  $n$ -dimensional  $k$ - $IC^\bullet$  manifold.*

*Then, every open subset of  $X$  is a  $k$ - $IC^\bullet$  manifold,  $X$  is purely  $n$ -dimensional, and the irreducible components of  $X$  are disjoint from each other.*

*Thus, for each irreducible component  $X_0$  of  $X$ ,  $k_{X_0}^\bullet[n]$  is a simple object in  $\text{Perv}(X_0)$ .*

*Proof.* Conditions 3), 5), 6), and 7) of Theorem 1.1 are clearly local; hence, every open subset of a  $k$ - $IC^\bullet$  manifold is a  $k$ - $IC^\bullet$  manifold.

If  $X$  had a component,  $X_0$ , of dimension less than  $n$ , then at a generic (smooth) point of  $X_0$ , condition 5) of Theorem 1.1 would fail.

Let  $X_1$  be an irreducible component of  $X$  such that  $X_1 \cap \overline{X - X_1} \neq \emptyset$ . Then there is a Whitney stratification of  $X$  such that  $X_1 \cap \overline{X - X_1}$  is a union of strata. Let  $S$  be a top-dimensional stratum of  $X_1 \cap \overline{X - X_1}$ . Then, the real link of  $S$  has more than one connected component; this would contradict 7) from Theorem 1.1.  $\square$

**Corollary 1.4.** *Let  $X$  and  $Y$  be arbitrary analytic spaces. Then,  $X \times Y$  is a  $k$ - $IC^\bullet$  manifold if and only if  $X$  and  $Y$  are both  $k$ - $IC^\bullet$  manifolds.*

*Proof.* We believe that it is well-known that the product of homology manifolds is a homology manifold; however, for lack of a convenient reference, we supply a proof. The technical results that we contained in Corollary 2.0.4 of [Sch].

Let  $n$  and  $m$  denote the (global) dimensions of  $X$  and  $Y$ , respectively. Let  $\pi_X$  and  $\pi_Y$  denote the projections from  $X \times Y$  onto  $X$  and  $Y$ , respectively. Then,

$$\mathcal{D}(k_{X \times Y}^\bullet[n + m]) \cong \pi_X^* \mathcal{D}(k_X^\bullet[n]) \overset{L}{\otimes} \pi_Y^* \mathcal{D}(k_Y^\bullet[m]),$$

and so, for all  $(p, q) \in X \times Y$ , for all  $i$ ,

$$H^i(\mathcal{D}(k_{X \times Y}^\bullet[n + m]))_{(p, q)} \cong \bigoplus_{s+t=i} H^s(\mathcal{D}(k_X^\bullet[n]))_p \otimes H^t(\mathcal{D}(k_Y^\bullet[m]))_q.$$

Using 3) from Theorem 1.1, it follows immediately that if  $X$  and  $Y$  are  $k$ - $IC^\bullet$  manifolds, then so is  $X \times Y$ .

Now suppose that  $X \times Y$  is a  $k$ - $IC^\bullet$  manifold. By selecting  $p$  to be a smooth point on an  $n$ -dimensional component of  $X$ , and allowing  $q$  to vary over  $Y$ , we find (again using 3) from Theorem 1.1) that  $Y$  must be a  $k$ - $IC^\bullet$  manifold; the analogous argument implies that  $X$  must be a  $k$ - $IC^\bullet$  manifold.  $\square$

*Remark 1.5.* Since being a  $k$ -intersection cohomology manifold is a local property, Corollary 1.4 immediately implies that an analytic space which fibers locally trivially over a  $k$ - $IC^\bullet$  manifold, with a fiber which is also a  $k$ - $IC^\bullet$  manifold, must itself be a  $k$ - $IC^\bullet$  manifold.

## §2. Hypersurfaces and Vanishing Cycles

Recall that  $f : X \rightarrow \mathbb{C}$  is a complex analytic function. Let  $j$  denote the closed inclusion of  $V(f)$  into  $X$ , and let  $i$  denote the open inclusion of  $X - V(f)$  into  $X$ . We must recall a number of basic results in the derived category  $D_c^b(X)$ .

The composed functors  $Ri_*i^*$  and  $i_1i^!$  take perverse sheaves to perverse sheaves. The shifted nearby cycle functor  $\psi_f[-1]$  is a functor from  $D_c^b(X)$  to  $D_c^b(V(f))$  which takes perverse sheaves to perverse sheaves. The nearby cycle monodromy  $T_f$  is a natural automorphism of the functor  $\psi_f[-1]$ . The shifted vanishing cycle functor  $\phi_f[-1]$  is a functor from  $D_c^b(X)$  to  $D_c^b(V(f))$  which takes perverse sheaves to perverse sheaves. The vanishing cycle monodromy  $\tilde{T}_f$  is a natural automorphism of the functor  $\phi_f[-1]$ .

When the complex of sheaves,  $\mathbf{A}^\bullet$ , under consideration is clear, we normally continue to write  $T_f$  and  $\tilde{T}_f$  for the morphisms  $T_f(\mathbf{A}^\bullet) : \psi_f[-1]\mathbf{A}^\bullet \rightarrow \psi_f[-1]\mathbf{A}^\bullet$  and  $\tilde{T}_f(\mathbf{A}^\bullet) : \phi_f[-1]\mathbf{A}^\bullet \rightarrow \phi_f[-1]\mathbf{A}^\bullet$ .

Let  $\mathbf{A}^\bullet \in D_c^b(X)$ . There are four canonical distinguished triangles that we shall need:

$$(i) \quad \mathbf{A}^\bullet \rightarrow Ri_*i^*\mathbf{A}^\bullet \rightarrow j_!j^![1]\mathbf{A}^\bullet \xrightarrow{[1]} ;$$

$$(ii) \quad j_*j^*[-1]\mathbf{A}^\bullet \rightarrow i_1i^!\mathbf{A}^\bullet \rightarrow \mathbf{A}^\bullet \xrightarrow{[1]} ;$$

$$(iii) \quad j^*[-1]\mathbf{A}^\bullet \xrightarrow{\text{comp}} \psi_f[-1]\mathbf{A}^\bullet \xrightarrow{\text{can}} \phi_f[-1]\mathbf{A}^\bullet \xrightarrow{[1]} ;$$

and

$$(iv) \quad \phi_f[-1]\mathbf{A}^\bullet \xrightarrow{\text{var}} \psi_f[-1]\mathbf{A}^\bullet \xrightarrow{\text{pmoc}} j^![1]\mathbf{A}^\bullet \xrightarrow{[1]} .$$

There are equalities  $\text{var} \circ \text{can} = \text{id} - T_f$  and  $\text{can} \circ \text{var} = \text{id} - \tilde{T}_f$ . We let  $\omega_f$  denote the morphism from  $j^*[-1]\mathbf{A}^\bullet$  to  $j^![1]\mathbf{A}^\bullet$  given by  $\text{pmoc} \circ \text{comp}$ . The octahedral lemma implies that the mapping cones of  $\omega_f$  and  $\text{id} - T_f$  are isomorphic; in particular,  $\omega_f$  is an isomorphism if and only if  $\text{id} - T_f$  is an isomorphism. We refer to  $\omega_f$  as the *Wang morphism*.

The triples  $(\text{id}, T_f, \tilde{T}_f)$  and  $(\tilde{T}_f, T_f, \text{id})$  yield automorphisms of the distinguished triangles in (iii) and (iv), respectively. In particular, if  $f$  is identically zero on  $X$ , then  $\psi_f[-1]\mathbf{A}^\bullet = 0$  and so  $\phi_f[-1]\mathbf{A}^\bullet \cong j^*\mathbf{A}^\bullet \cong j^!\mathbf{A}^\bullet$ , and  $\tilde{T}_f$  is the identity.

Suppose now that  $\mathbf{A}^\bullet$  is a simple object in  $\text{Perv}(X)$ . If  $\text{supp } \mathbf{A}^\bullet \not\subseteq V(f)$ , then the distinguished triangles (i) and (ii) imply that  $j_!j^![1]\mathbf{A}^\bullet$  and  $j_*j^*[-1]\mathbf{A}^\bullet$  are in  $\text{Perv}(X)$  and, thus,  $j^![1]\mathbf{A}^\bullet$  and  $j^*[-1]\mathbf{A}^\bullet$  are in  $\text{Perv}(V(f))$ .

**Lemma 2.1.** *Suppose that  $X$  is an  $n$ -dimensional  $k$ -IC $^\bullet$  manifold.*

*If  $f$  does not vanish identically on any irreducible component of  $X$ , then  $j^*[-1]k_x^\bullet[n]$  and  $j^![1]k_x^\bullet[n]$  are perverse sheaves on  $V(f)$ , which are Verdier dual to each other, and the image (in  $\text{Perv}(V(f))$ ) of  $\text{id} - T_f : \psi_f[-1]k_x^\bullet[n] \rightarrow \psi_f[-1]k_x^\bullet[n]$  is isomorphic to  $\phi_f[-1]k_x^\bullet[n]$ .*

*If  $f$  vanishes identically on an irreducible component  $X_0$  of  $X$ , then, when restricted to  $X_0$ ,  $\text{id} - \tilde{T}_f : \phi_f[-1]k_x^\bullet[n] \rightarrow \phi_f[-1]k_x^\bullet[n]$  is the zero morphism and, hence, does not induce an isomorphism in any stalk at a point of  $X_0$ .*

*Proof.* Suppose that  $f$  does not vanish identically on any irreducible component of  $X$ . As the restriction of  $k_x^\bullet[n]$  to each irreducible component of  $X$  is a simple object, the perversity of  $j^*[-1]k_x^\bullet[n]$

and  $j^1[1]k_x^\bullet[n]$  follows from the case discussed above of a simple perverse sheaf. As  $X$  is a  $k$ -IC $^\bullet$  manifold,  $\mathcal{D}(k_x^\bullet[n]) \cong k_x^\bullet[n]$ , and so  $\mathcal{D}(j^*[-1]k_x^\bullet[n]) \cong j^1[1]\mathcal{D}(k_x^\bullet[n]) \cong j^1[1]k_x^\bullet[n]$ . Since  $j^*[-1]k_x^\bullet[n]$  and  $j^1[1]k_x^\bullet[n]$  are perverse sheaves, the distinguished triangles (iii) and (iv) above become short exact sequences in  $\text{Perv}(V(f))$  when we replace  $\mathbf{A}^\bullet$  with  $k_x^\bullet[n]$ . Thus, the morphism  $\text{can}$  is a surjection and the morphism  $\text{var}$  is an injection. It follows that the image of  $\text{id} - T_f = \text{var} \circ \text{can}$  is isomorphic to  $\phi_f[-1]k_x^\bullet[n]$ . This proves the first statement.

Suppose that  $f$  vanishes identically on an irreducible component  $X_0$  of  $X$ . From the paragraphs preceding the lemma, we know that, when restricted to  $X_0$ ,  $\phi_f[-1]k_x^\bullet[n] \cong k_x^\bullet[n]$  and that  $\text{id} - \widetilde{T}_f$  is the zero morphism. This proves the second statement.  $\square$

Now we come to our main theorem, which relates the vanishing cycle monodromy to  $k$ -intersection cohomology manifolds.

**Theorem 2.2.** *Let  $X$  be an  $n$ -dimensional  $k$ -IC $^\bullet$  manifold. The following are equivalent:*

- a)  $\text{id} - \widetilde{T}_f : \phi_f[-1]k_x^\bullet[n] \rightarrow \phi_f[-1]k_x^\bullet[n]$  is an isomorphism;
- b)  $V(f)$  is a  $k$ -IC $^\bullet$  manifold, and  $\text{id} - \widetilde{T}_f : \phi_f[-1]k_x^\bullet[n] \rightarrow \phi_f[-1]k_x^\bullet[n]$  is an isomorphism when restricted to a generic subset of  $V(f)$ ;
- c) the pairs of complexes and morphisms  $(\psi_f[-1]k_x^\bullet[n]; T_f)$  and  $(k_{V(f)}^\bullet[n-1] \oplus \phi_f[-1]k_x^\bullet[n]; \text{id} \oplus \widetilde{T}_f)$  are isomorphic in  $D_c^b(V(f))$ , i.e., there is an isomorphism  $\theta : \psi_f[-1]k_x^\bullet[n] \rightarrow k_{V(f)}^\bullet[n-1] \oplus \phi_f[-1]k_x^\bullet[n]$  such that  $\theta \circ T_f = (\text{id} \oplus \widetilde{T}_f) \circ \theta$ ;

and, in these equivalent cases,  $V(f)$  contains no component of  $X$ ,  $V(f)$  is purely  $(n-1)$ -dimensional, and the isomorphism in c) is an isomorphism in  $\text{Perv}(V(f))$ .

*Proof.* By Lemma 2.1, a) and b) both imply that  $V(f)$  contains no component of  $X$ . If  $\psi_f[-1]k_x^\bullet[n] \cong k_{V(f)}^\bullet[n-1] \oplus \phi_f[-1]k_x^\bullet[n]$ , then, since  $\psi_f[-1]k_x^\bullet[n]$  and  $\phi_f[-1]k_x^\bullet[n]$  are perverse, it follows that  $k_{V(f)}^\bullet[n-1]$  is perverse; as  $k_x^\bullet[n]$  is perverse when restricted to each irreducible component, this implies that  $V(f)$  cannot contain a component of  $X$ . Therefore, by Lemma 2.1, a), b), and c) all imply that  $V(f)$  contains no component of  $X$ ,  $V(f)$  is purely  $(n-1)$ -dimensional, and  $j^*[-1]k_x^\bullet[n]$  and  $j^1[1]k_x^\bullet[n]$  are perverse sheaves on  $V(f)$ , which are Verdier dual to each other. In particular, if the isomorphism in c) holds, then it is an isomorphism in  $\text{Perv}(V(f))$ .

Proof of a)  $\implies$  b): Suppose that  $\text{id} - \widetilde{T}_f$  is an isomorphism.

As  $\text{id} - \widetilde{T}_f$  is an isomorphism, the Wang morphism  $\omega_f : j^*[-1]k_x^\bullet[n] \rightarrow j^1[1]k_x^\bullet[n]$  is an isomorphism. Thus,

$$k_{V(f)}^\bullet[n-1] \cong j^*[-1]k_x^\bullet[n] \cong j^1[1]k_x^\bullet[n] \cong \mathcal{D}(j^*[-1]k_x^\bullet[n]) \cong \mathcal{D}(k_{V(f)}^\bullet[n-1]).$$

By 2) of Theorem 1.1,  $V(f)$  is an  $(n-1)$ -dimensional  $k$ -IC $^\bullet$  manifold.

Proof of b)  $\implies$  a): Suppose that  $\text{id} - \widetilde{T}_f : \phi_f[-1]k_x^\bullet[n] \rightarrow \phi_f[-1]k_x^\bullet[n]$  is an isomorphism when restricted to a generic subset of  $V(f)$ , and  $V(f)$  is a  $k$ -IC $^\bullet$  manifold.

Since  $\text{id} - \tilde{T}_f$  is an isomorphism on a generic subset of  $V(f)$ , the Wang morphism  $\omega_f : j^*[-1]k_x^\bullet[n] \rightarrow j^![-1]k_x^\bullet[n]$  is an isomorphism on a generic subset of  $V(f)$ . As  $V(f)$  is an  $(n-1)$ -dimensional  $k\text{-IC}^\bullet$  manifold,  $k_{V(f)}^\bullet[n-1] \cong j^*[-1]k_x^\bullet[n]$  is simple when restricted to each irreducible component of  $V(f)$ . As  $j^![-1]k_x^\bullet[n] \cong \mathcal{D}(j^*[-1]k_x^\bullet[n])$ ,  $j^![-1]k_x^\bullet[n]$  is also simple when restricted to each irreducible component of  $V(f)$ . Thus,  $\omega_f$  must be an isomorphism on all of  $V(f)$ , and, hence, so is  $\text{id} - \tilde{T}_f$ .

Proof of a)  $\implies$  c): Assume a). As  $j^*[-1]k_x^\bullet[n]$  is perverse, the canonical distinguished triangle (iii) from the beginning of the section becomes a short exact sequence in the category of perverse sheaves on  $V(f)$

$$(†) \quad 0 \rightarrow k_{V(f)}^\bullet[n-1] \xrightarrow{\text{comp}} \psi_f[-1]k_x^\bullet[n] \xrightarrow{\text{can}} \phi_f[-1]k_x^\bullet[n] \rightarrow 0.$$

Recall that the variation natural transformation yields a morphism  $\text{var} : \phi_f[-1]k_{\mathcal{U}}^\bullet[n] \rightarrow \psi_f[-1]k_{\mathcal{U}}^\bullet[n]$ , and  $\text{can} \circ \text{var} = \text{id} - \tilde{T}_f$ . Thus,  $\text{var} \circ (\text{id} - \tilde{T}_f)^{-1}$  yields a splitting of (†). Moreover, this splitting is compatible with the morphism of (†) given by the triple  $(\text{id}, T_f, \tilde{T}_f)$ . The isomorphism of complexes and automorphisms in c) follows.

Proof of c)  $\implies$  a): Assume c). As  $V(f)$  contains no component of  $X$ , the first statement of Lemma 2.1 implies that  $\text{im}(\text{id} - T_f) \cong \phi_f[-1]k_x^\bullet[n]$ . On the other hand, the isomorphism of the complexes and automorphisms implies that  $\text{im}(\text{id} - T_f) \cong 0 \oplus \text{im}(\text{id} - \tilde{T}_f)$ . Therefore,  $\text{im}(\text{id} - \tilde{T}_f) \cong \phi_f[-1]k_x^\bullet[n]$ . As  $\text{Perv}(V(f))$  is locally Artinian, it follows that  $\text{id} - \tilde{T}_f$  is an isomorphism.  $\square$

*Remark 2.3.* The condition in Theorem 2.2.b that  $\text{id} - \tilde{T}_f$  is an isomorphism when restricted to a generic subset of  $V(f)$  may seem like a very strong, unusual condition. In fact, this condition is what one expects to happen, for on the open set  $\mathcal{U} := V(f) - \text{supp}(\phi_f[-1]k_x^\bullet[n])$ ,  $\phi_f[-1]k_x^\bullet[n]$  is zero and thus  $(\text{id} - \tilde{T}_f)|_{\mathcal{U}}$  is an isomorphism.

Consider, for example, the case where  $X$  is an open subset of  $\mathbb{C}^n$ . If  $f$  is reduced, then  $\phi_f[-1]k_x^\bullet[n]$  is zero on a generic subset of  $V(f)$ , and so  $\text{id} - \tilde{T}_f$  is an isomorphism on a generic subset of  $V(f)$ . In this case, one can prove the equivalence of a) and b) in Theorem 2.2 via the Wang sequence, precisely as Milnor does in the isolated case in Chapter 8 of [Mi].

Suppose now that  $f$  is not reduced, and  $f = f_1^{\alpha_1} f_2^{\alpha_2} \dots f_k^{\alpha_k}$  is the irreducible factorization of  $f$  at a point  $\mathbf{p} \in X$ . At a generic point  $\mathbf{q}_i \in V(f_i)$ , up to isomorphism, we have the case where  $f$  is a power of a coordinate function, i.e.,  $f = z_0^{\alpha_i}$ . By taking a normal slice, we are reduced to considering the case where  $f : \mathbb{C} \rightarrow \mathbb{C}$  is given by  $f(z) = z^{\alpha_i}$ . The Milnor fiber consists of  $\alpha_i$  points and the monodromy  $T_f : k^{\alpha_i} \rightarrow k^{\alpha_i}$  is simply a cyclic permutation of the coordinates. Let  $\Delta$  denote the diagonal in  $k^{\alpha_i}$ . The map induced by  $\text{id} - \tilde{T}_f$  is simply the automorphism of  $k^{\alpha_i}/\Delta$  induced by  $\text{id} - T_f$ . It is trivial to show that this map is an isomorphism if and only if  $\alpha_i$  is not divisible by the characteristic of  $k$ . Therefore, we conclude that if the characteristic of  $k$  does not divide the multiplicity of any component of  $f$ , then  $\text{id} - \tilde{T}_f$  is an isomorphism on a generic subset of  $V(f)$ .

*Remark 2.4.* In the case where  $X$  is smooth,  $f$  is reduced, and  $k = \mathbb{C}$ , Tristan Torrelli [T] uses the language and techniques of  $\mathcal{D}$ -modules to establish the same equivalence as we obtain in a) and b) of our Theorem 2.2. Moreover, Torrelli includes one more condition which is equivalent to  $V(f)$  being a  $\mathbb{C}\text{-IC}^\bullet$  manifold: namely, that the reduced Bernstein polynomial of  $f$  has no integral root.

The following topological result was suggested to us by Alberto Verjovsky. It says that in the classical affine situation, if the singular set of the hypersurface  $V(f)$  is not too big, then we have a fairly obvious generalization of the results of Milnor in Chapter 8 of [Mi].

**Theorem 2.5.** *Let  $X$  be an open subset of  $\mathbb{C}^n$ . Let  $s$  denote the dimension of the singular set  $\Sigma V(f)$ , and suppose that  $n - 1 - s \geq 3$ . Then, the following are equivalent:*

- 1)  $V(f)$  is an integral homology (or cohomology) manifold;
- 2)  $V(f)$  is a  $k$ -IC<sup>•</sup> manifold for  $k = \mathbb{Q}$  (or  $\mathbb{C}$ ) and for  $k = \mathbb{Z}/p\mathbb{Z}$  for all primes  $p$ ;
- 3) for all complex analytic Whitney stratifications of  $V(f)$ , the real links of all strata are topological spheres;
- 4) there exists a complex analytic Whitney stratification of  $V(f)$  such that the real links of all strata are topological spheres;
- 5)  $V(f)$  is a topological manifold.

*Proof.* For all  $x \in X$ , the real link of  $X$  at  $x$  has the homotopy-type of a finite CW-complex and, hence, the integral homology groups are finitely-generated. Therefore, the equivalence of 1) and 2) follow from the Universal Coefficient Theorems, together with Theorem 1.1.

Certainly, 3) implies 4), and 5) implies 1).

Suppose that  $S$  is a Whitney stratum of  $X$ . Then, along  $S$ ,  $X$  is locally a product of Euclidean space with the cone on the real link of  $S$ . Hence, 4) implies 5).

We need to prove that 1) implies 3).

Assume that  $V(f)$  is an integral homology manifold, and let  $\mathcal{S}$  be a Whitney stratification of  $V(f)$ . By the equivalence of 1) and 2), combined with Theorem 1.1, we find that the real link of each stratum of  $\mathcal{S}$  is an integral homology sphere.

The real link of a stratum is the real link of a point in the normal slice. As  $n - 1 - s \geq 3$ , when we take the real link of a point in a normal slice to a singular stratum, that link will be at least 5-dimensional and will be simply-connected by Theorem 5.2 of [Mi].

Therefore, the real links of the strata are at least 5-dimensional and are homotopy-equivalent to spheres. If we can show that the real links to strata are topological manifolds, then we will be finished by the Generalized Poincaré Conjecture (Theorem).

We prove this by contradiction. Let  $S \in \mathcal{S}$  be a stratum of maximum dimension such that its real link,  $K$ , is not a topological manifold. The link  $K$  is the transverse intersection of a small sphere in the ambient space with a normal slice to  $V(f)$ . This link inherits a Whitney stratification  $\mathcal{S}'$  from  $\mathcal{S}$ . The links of strata of  $\mathcal{S}'$  are precisely the links of the corresponding strata of  $\mathcal{S}$ ; by our maximality assumption on  $S$ , the links of the strata of  $\mathcal{S}'$  are topological spheres. The topology of  $K$  along a stratum of  $\mathcal{S}'$  is a product of Euclidean space with the cone on a sphere. Thus,  $K$  is a manifold; a contradiction.  $\square$

### §3. Suspending $f$

For  $j \in \mathbb{N}$ ,  $j \geq 2$ , let  $f_j : X \times \mathbb{C} \rightarrow \mathbb{C}$  be given by  $f_j(\mathbf{z}, w) := f(\mathbf{z}) + w^j$ . We refer to  $f_j$  as the  $(j - 1)$ -fold suspension of  $f$ . We use this terminology because the homotopy-type of the Milnor fiber

of  $f_j$  at a point of  $(\mathbf{p}, 0) \in V(f) \times \{0\}$  is the one-point union of  $(j-1)$  copies of the suspension of the Milnor fiber of  $f$  at  $\mathbf{p}$ ; see the references below.

Our interest in suspending  $f$  stems from the Sebastiani-Thom isomorphism (see [O], [Sa], [S-T], [N1], [N2], [Sch], [Mas4]), which implies that  $\text{supp}(\phi_{f_j}[-1]k_{x \times \mathbb{C}}^\bullet[n+1]) = \text{supp}(\phi_f[-1]k_x^\bullet[n]) \times \{0\}$  and, at a point  $(\mathbf{p}, 0) \in V(f) \times \{0\}$ ,

$$H^i(\phi_{f_j}[-1]k_{x \times \mathbb{C}}^\bullet[n+1])_{(\mathbf{p}, 0)} \cong H^i(\phi_f[-1]k_x^\bullet[n])_{\mathbf{p}} \otimes k^{j-1},$$

i.e.,

$$\tilde{H}^{n+i}(F_{f_j, (\mathbf{p}, 0)}) \cong (\tilde{H}^{n+i-1}(F_{f, \mathbf{p}}))^{j-1},$$

and that the action of  $\tilde{T}_{f_j}$  on the stalk cohomology of the vanishing cycles of  $f_j$  is the tensor product of the action of  $\tilde{T}_f$  on the stalk cohomology of the vanishing cycles of  $f$  at  $\mathbf{p}$  and the action on  $k^{j-1}$  given by a cyclically permuting the coordinates of  $k^j$  and modding out by the diagonal.

**Definition 3.1.** A point  $\mathbf{p} \in V(f)$  is *totally  $f$ -unipotent* if and only if, at every point  $\mathbf{q} \in V(f)$  near  $\mathbf{p}$ , and in all degrees  $i$ , if  $H^i(\phi_f[-1]k_x^\bullet[n])_{\mathbf{p}} \neq 0$ , then the characteristic polynomial,  $\text{char}_{(\tilde{T}_f^i)_{\mathbf{q}}}(t)$ , of the stalk of  $\tilde{T}_f$  at  $\mathbf{q}$  in degree  $i$  is a power of  $(t-1)$ .

Note that if  $\mathbf{p} \in V(f) - \text{supp}(\phi_f[-1]k_x^\bullet[n])$ , then  $\mathbf{p}$  is totally  $f$ -unipotent.

**Theorem 3.2.** *Let  $X$  be an  $n$ -dimensional  $k$ -IC $^\bullet$  manifold. Suppose that  $f$  does not vanish identically on a component of  $X$ . Let  $\mathbf{p} \in V(f)$ . Let*

$$J := \{j \in \mathbb{N} \mid j \geq 2, V(f_j) \text{ is a } k\text{-IC}^\bullet \text{ manifold in a neighborhood of } (\mathbf{p}, 0)\}.$$

*Then,  $J$  is an infinite set. Moreover, if  $\mathbf{p}$  is totally  $f$ -unipotent, then  $J = \{j \in \mathbb{N} \mid j \geq 2\}$ .*

*In addition, if  $\mathbf{p}$  is not totally  $f$ -unipotent, then  $\mathbb{N} - J$  is also an infinite set.*

*Proof.* First note that Corollary 1.4 implies that  $X \times \mathbb{C}$  is a  $k$ -IC $^\bullet$  manifold.

Now, as  $\text{supp}(\phi_{f_j}[-1]k_{x \times \mathbb{C}}^\bullet[n+1]) = \text{supp}(\phi_f[-1]k_x^\bullet[n]) \times \{0\}$  and since we are assuming that  $f$  does not vanish identically on a component of  $X$ , the dimension of the support of  $\phi_{f_j}[-1]k_{x \times \mathbb{C}}^\bullet[n+1]$  is, at most,  $n-1$ . However, the dimension of each component of  $V(f_j)$  is precisely  $n$ . Therefore,  $\text{id} - \tilde{T}_{f_j}$  is an isomorphism when restricted to a generic subset of  $V(f_j)$ , and Theorem 2.2 implies that  $V(f_j)$  is a  $k$ -IC $^\bullet$  manifold if and only if  $\text{id} - \tilde{T}_{f_j}$  is an isomorphism on all of  $V(f_j)$ .

Since 1 is an eigenvalue of a linear operator over  $k$  if and only if 1 is an eigenvalue of the induced linear operator over some extension of  $k$ , we may assume that  $k$  is algebraically closed.

Fix a Whitney stratification of  $X$  such that  $V(f)$  is a union of strata. Define a  $\mathbf{p}$ -stratum to be a stratum of this Whitney stratification which contains  $\mathbf{p}$  in its closure. Let  $D$  be the set of  $v \in \mathbb{C}$  such that there exists  $x \in \text{supp} \phi_f[-1]k_x^\bullet[n]$  such that  $x$  is in a  $\mathbf{p}$ -stratum and  $v$  is an eigenvalue of the stalk at  $x$  of the vanishing cycle monodromy in some degree. Then, by constructibility,  $D$  is a finite set, and by the Monodromy Theorem (see [Cl], [Gr], [La], [Lê2])  $D$  consists of roots of unity. Let  $E$  be the set of natural numbers  $q$  such that a primitive  $q$ -th root of unity is contained in  $D$ . Then,  $E$  is a finite set.

Note that  $\mathbf{p}$  is totally  $f$ -unipotent if and only if  $E = \{1\}$  or  $E = \emptyset$ .

If  $E = \emptyset$ , then  $\mathbf{p} \notin \text{supp}(\phi_f[-1]k_x^\bullet[n])$ . Thus, for all  $j$ ,  $(\mathbf{p}, 0) \notin \text{supp}(\phi_{f_j}[-1]k_{x \times \mathbb{C}}^\bullet[n+1])$ , and so, in a neighborhood of  $(\mathbf{p}, 0)$ ,  $\text{id} - \tilde{T}_{f_j}$  induces isomorphisms on stalk cohomology (since the stalk cohomology is zero). Therefore, Theorem 2.2 implies that  $J = \{j \in \mathbb{N} \mid j \geq 2\}$ .

Suppose now that  $E \neq \emptyset$ .

By the Sebastiani-Thom Theorem, if  $j$  is relatively prime to each element of  $E$ , then the vanishing cycle monodromy of  $f_j$  does not have 1 as an eigenvalue in any stalk, and hence, by Theorem 2.2,  $j \in J$ .

On the other hand, if  $\mathbf{p}$  is not totally  $f$ -unipotent and  $j$  is a multiple of an element of  $E - \{1\}$ , then 1 will be an eigenvalue of the vanishing cycle monodromy  $\tilde{T}_{f_j}$  at points near  $(\mathbf{p}, 0)$ ; thus,  $V(f_j)$  is not a  $k\text{-IC}^\bullet$  manifold.  $\square$

The trick above of adding a power of  $w$  in order to eliminate 1 as an eigenvalue of the monodromy appears in [Lê1].

*Example 3.3.* Suspending functions is interesting even in the most basic case of the ordinary quadratic singularity  $f : \mathbb{C}^n \rightarrow \mathbb{C}$  given by  $f := z_1^2 + \cdots + z_n^2$ , which is an iterated suspension. The vanishing cycle monodromy is given by the single map multiplication by  $(-1)^n$  on  $\tilde{H}^{n-1}(F_{f, \mathbf{0}}) \cong k$ .

When  $n = 2$ , one can simply see the two irreducible components which prevent the shifted constant sheaf from being intersection cohomology. More generally, if  $n$  is even, the previous paragraph implies that  $V(f)$  is not a  $k\text{-IC}^\bullet$  manifold, regardless of what the field  $k$  is.

If  $n$  is odd, then  $V(f)$  is a  $k\text{-IC}^\bullet$  manifold, provided that  $k$  does not have characteristic 2.

If  $k$  has characteristic 2, then the ordinary quadratic singularity **never** yields a  $k\text{-IC}^\bullet$  manifold, except in the trivial case where  $n = 1$ .

The following is an easy corollary of Theorems 2.5 and 3.2.

**Corollary 3.4.** *Let  $X$  be an open subset of  $\mathbb{C}^n$ , and suppose that  $f = f(\mathbf{z})$  is reduced at each point of  $V(f)$ . If  $V(f)$  is an integral homology manifold, then the subset of  $X \times \mathbb{C}^2$  given by  $V(f(\mathbf{z}) + w_1^2 + w_2^2)$  is a topological manifold whose (analytic) singular set is isomorphic to the singular set of  $V(f)$ .*

*Proof.* Let  $s$  denote the dimension of the singular set  $\Sigma V(f)$ . As  $f$  is reduced,  $n - 1 - s \geq 1$ . Thus,  $(n + 2) - 1 - s \geq 3$ , and we are in a position to apply Theorem 2.5 to  $V(f(\mathbf{z}) + w_1^2 + w_2^2)$ .

Suppose  $V(f)$  is an integral homology manifold. As we saw in the proof of Theorem 2.5, this is equivalent to  $V(f)$  being a  $k\text{-IC}^\bullet$  manifold for  $k = \mathbb{Q}$  and  $k = \mathbb{Z}/p\mathbb{Z}$ , for all primes  $p$ . By the Sebastiani-Thom Theorem, at each point in the critical locus of  $f + w_1^2 + w_2^2$ , the eigenvalues of the vanishing cycle monodromy of  $f + w_1^2 + w_2^2$  are  $(-1)^2 = 1$  times the eigenvalues of the vanishing cycle monodromy of  $f$ . Thus, Theorem 2.2 and Remark 2.3 imply that  $V(f(\mathbf{z}) + w_1^2 + w_2^2)$  is an integral homology manifold. The desired result now follows from Theorem 2.5.  $\square$

The reader should compare the above “double suspension” result with that of Cannon in [Ca].

#### §4. One-Dimensional Critical Loci

For the sake of concreteness and ease of notation, we assume throughout this section that  $X = \mathcal{U}$  is a connected open neighborhood of the origin in  $\mathbb{C}^n$ , that  $f : \mathcal{U} \rightarrow \mathbb{C}$  has a one-dimensional critical locus through the origin, and that  $n \geq 3$ . Then,  $f$  is reduced at the origin, and the only possibly non-zero reduced cohomology vector spaces of  $F_{f, \mathbf{0}}$  occur in degrees  $n - 2$  and  $n - 1$ .

Select a linear form  $z_0$  so that  $f|_{V(z_0)}$  has an isolated critical point at  $\mathbf{0}$ , and define a family of isolated singularities  $f_t := f|_{V(z_0 - t)}$  (throughout this section, we assume that our neighborhood  $\mathcal{U}$  is re-chosen as small as necessary). Fix a small  $t_0 \neq 0$ . In this section, we will show that if  $V(f_{t_0})$  is a  $k$ -IC $^\bullet$  manifold and  $V(f_0)$  is not, then there are restrictions on  $\tilde{H}^{n-2}(F_{f, \mathbf{0}})$  (and, hence, on  $\tilde{H}^{n-1}(F_{f, \mathbf{0}})$ ). Note that  $f_0$  and  $f_{t_0}$  are both reduced near the origin.

First we need to recall some fairly well-known background material.

Let  $\mathbf{P}^\bullet$  denote the perverse sheaf  $(\phi_f[-1]k_{\mathcal{U}}^\bullet[n])|_{\Sigma f}$ . Assume we are in a small enough neighborhood of the origin so that  $\mathbf{P}^\bullet$  is constructible with respect to  $\{\Sigma f - \{\mathbf{0}\}, \{\mathbf{0}\}\}$ . Let  $\nu$  be an irreducible component of  $\Sigma f$ . Let  $\mathbf{p}_\nu$  denote a point on  $\nu - \{\mathbf{0}\}$ . Then,  $\mathbf{P}^\bullet|_{\nu - \mathbf{0}}[-1]$  is a local system with stalk isomorphic to  $\tilde{H}^{n-2}(F_{f_{t_0}, \mathbf{p}_\nu}) \cong k^{\mu_\nu}$ , where  $\mu_\nu$  is the “generic Milnor number along  $\nu$ ” near  $\mathbf{0}$ . As  $\nu - \mathbf{0}$  is homotopy-equivalent to a circle,  $\mathbf{P}^\bullet|_{\nu - \mathbf{0}}[-1]$  is characterized by a monodromy automorphism  $h_\nu$  on  $\tilde{H}^{n-2}(F_{f_{t_0}, \mathbf{p}_\nu})$ . As  $\tilde{T}_f$  induces an automorphism on the complex  $\mathbf{P}^\bullet$ , the Milnor monodromy induced on each  $\tilde{H}^{n-2}(F_{f_{t_0}, \mathbf{p}_\nu})$  must commute with  $h_\nu$ .

Let  $\hat{m} : \{\mathbf{0}\} \hookrightarrow \Sigma f$  denote the closed inclusion, and let  $\hat{l} : \Sigma f - \{\mathbf{0}\} \hookrightarrow \Sigma f$  denote the open inclusion. There is a canonical distinguished triangle (see (i) at the beginning of Section 2)

$$(\dagger) \quad \hat{m}_! \hat{m}^! \mathbf{P}^\bullet \rightarrow \mathbf{P}^\bullet \rightarrow R\hat{l}_* \hat{l}^* \mathbf{P}^\bullet \rightarrow \hat{m}_! \hat{m}^! \mathbf{P}^\bullet[1].$$

The cosupport condition on the perverse sheaf  $\mathbf{P}^\bullet$  implies that  $H^{-1}(\mathbf{P}^\bullet)_\mathbf{0} = 0$ , and so the long exact sequence on stalk cohomology obtained from  $(\dagger)$  yields that there is an injection  $\beta$  given by the following composition

$$\beta : \tilde{H}^{n-2}(F_{f, \mathbf{0}}) \cong H^{-1}(\mathbf{P}^\bullet)_\mathbf{0} \hookrightarrow H^{-1}(R\hat{l}_* \hat{l}^* \mathbf{P}^\bullet)_\mathbf{0} \cong \bigoplus_\nu \ker(\text{id} - h_\nu) \hookrightarrow \bigoplus_\nu \tilde{H}^{n-2}(F_{f_{t_0}, \mathbf{p}_\nu}),$$

where  $\beta$  commutes with the respective Milnor monodromies. This injection was first proved by Siersma in [Si].

**Theorem 4.1.** *The vector space  $\tilde{H}^{n-2}(F_{f, \mathbf{0}})$  injects into both  $\tilde{H}^{n-2}(F_{f_0, \mathbf{0}})$  and  $\bigoplus_\nu \tilde{H}^{n-2}(F_{f_{t_0}, \mathbf{p}_\nu})$  by inclusions which commute with the actions of the respective Milnor monodromies.*

*If  $\dim \tilde{H}^{n-2}(F_{f, \mathbf{0}}) = \sum_\nu \mu_\nu$ , then each  $h_\nu$  is the identity, and the eigenspaces of the Milnor monodromy on  $\bigoplus_\nu \tilde{H}^{n-2}(F_{f_{t_0}, \mathbf{p}_\nu})$  inject into the corresponding eigenspaces of the Milnor monodromy on  $\tilde{H}^{n-2}(F_{f_0, \mathbf{0}})$ .*

*Proof.* The discussion above, especially the definition of the injection  $\beta$ , tells us that  $\tilde{H}^{n-2}(F_{f, \mathbf{0}})$  injects into  $\bigoplus_\nu \tilde{H}^{n-2}(F_{f_{t_0}, \mathbf{p}_\nu})$  by an inclusion which commutes with the actions of the respective Milnor monodromies.

The Main Theorem of [Mas2] implies that  $\tilde{H}^{n-2}(F_{f,\mathbf{0}})$  injects into  $\tilde{H}^{n-2}(F_{f_0,\mathbf{0}})$  by an inclusion which commutes with the actions of the respective Milnor monodromies.

If  $\dim \tilde{H}^{n-2}(F_{f,\mathbf{0}}) = \sum_{\nu} \overset{\circ}{\mu}_{\nu}$ , then  $\beta$  must be an isomorphism. Hence, the second statement follows immediately from the first statement.  $\square$

**Corollary 4.2.** *If  $V(f_{t_0})$  is not a  $k\text{-IC}^{\bullet}$  manifold, and  $V(f_0)$  is, then  $\dim \tilde{H}^{n-2}(F_{f,\mathbf{0}}) < \sum_{\nu} \overset{\circ}{\mu}_{\nu}$ .*

*Proof.* If  $V(f_{t_0})$  is not a  $k\text{-IC}^{\bullet}$  manifold, then, by Theorem 2.2, the Milnor monodromy on at least one of the  $\tilde{H}^{n-2}(F_{f_{t_0},\mathbf{p}_{\nu}})$  must have 1 as an eigenvalue. However, if  $V(f_0)$  is a  $k\text{-IC}^{\bullet}$  manifold, then Theorem 2.2 implies that 1 is not an eigenvalue of the Milnor monodromy on  $\tilde{H}^{n-2}(F_{f_0,\mathbf{0}})$ . It follows at once from Theorem 4.1 that  $\dim \tilde{H}^{n-2}(F_{f,\mathbf{0}}) < \sum_{\nu} \overset{\circ}{\mu}_{\nu}$ .  $\square$

### §5. Questions and Remarks

Corollary 4.2 is an interesting little result. It leads one to ask the following more general question:

**Question.** If  $f$  has a one-dimensional critical locus at the origin, and  $\dim \tilde{H}^{n-2}(F_{f,\mathbf{0}}) = \sum_{\nu} \overset{\circ}{\mu}_{\nu}$ , does it follow that the critical locus is itself smooth and that  $f$  defines a family of isolated singularities with constant Milnor number?

In fact, the example of  $f := (y^2 - x^3)^2 + w^2$  shows us that, in general, the answer to the above question is “no”. However, if all of the components of the critical locus of  $f$  are smooth at the origin, then we have showed in a recent paper [L-M], with Lê D. T., that the answer to the above question is “yes”.

We were trying to answer the above question when we proved most of the results of this paper. We hoped that, by suspending  $f$ , we could reduce ourselves to the case where  $V(f)$  is an intersection cohomology manifold, and that we could then make effective use of the nice categorical property of intersection cohomology: namely, that the simple objects in the locally Artinian category of perverse sheaves are precisely the intersection cohomology complexes on irreducible subvarieties with coefficients in irreducible local systems.

One might also hope to gain more information by considering the nearby cycles  $\psi_f[-1]k_{\mathcal{U}}^{\bullet}[n]$  instead of the vanishing cycles  $\phi_f[-1]k_{\mathcal{U}}^{\bullet}[n]$ . Theorem 2.2 tells us that the nearby cycles may contain interesting structure that is not present in the vanishing cycles precisely when  $V(f)$  is **not** an intersection cohomology manifold. Thus, it may be useful to suspend  $f$  in such a way that we do **not** have that  $V(f)$  is an intersection cohomology manifold; this is one of the reasons that we look at the set  $\mathbb{N} - J$  in Theorem 3.2.

Consider, for instance, a basic example which shows that the nearby cycles are “more interesting” than the vanishing cycles when  $V(f)$  is not an intersection cohomology manifold; let  $f : \mathbb{C}^2 \rightarrow \mathbb{C}$  be given by  $f(x, y) = x^2 + y^2$ . As we saw in Example 3.3, the monodromy map  $\tilde{T}_f$  on the vanishing cycles is simply the identity in this case. Does this imply that the monodromy map  $T_f$  is also the identity map? No. Recall that, in  $\text{Perv}(V(f))$ , the image of  $\text{id} - T_f$  is isomorphic to the vanishing cycles. Thus, if the vanishing cycles are non-zero, then  $T_f$  is not the identity. However, it is true that if  $\tilde{T}_f = \text{id}$ , then  $(\text{id} - T_f)^2 = 0$ .

## REFERENCES

- [A'C] N. A'Campo, *Le nombre de Lefschetz d'une monodromie*, Proc. Kon. Ned. Akad. Wet., Series A **76** (1973), 113–118.
- [B] D. Barlet, *Interaction de strates consécutives pour les cycles évanescents*, Ann. Sci. E.N.S. 4<sup>e</sup> série **24** (1991), 401–506.
- [B-M] W. Borho and R. MacPherson, *Partial Resolutions of Nilpotent Varieties*, Astérisque **101-102** (1982), 23–74.
- [Ca] J.W. Cannon, *The recognition problem: what is a topological manifold?*, Bull. AMS **84** (1978), 832–866.
- [Cl] C. H. Clemens, *Picard-Lefschetz theorem for families of non-singular algebraic varieties acquiring ordinary singularities*, Trans. AMS **136** (1969), 93–108.
- [D] A. Dimca, *Singularities and Topology of Hypersurfaces*, Universitext, Springer - Verlag, 1992.
- [G-M1] M. Goresky and R. MacPherson, *Morse Theory and Intersection Homology*, Analyse et Topologie sur les Espaces Singuliers. Astérisque **101** (1983), Soc. Math. France, 135–192.
- [G-M2] ———, *Intersection homology II*, Inv. Math **71** (1983), 77–129.
- [G-M3] ———, *Intersection homology theory*, Topology **19** (1980), 135–162.
- [Gr] A. Grothendieck, *Séminaire de Géométrie Algébrique (SGA VII-1), Résumé des premiers exposés de A. Grothendieck, rédigé par P. Deligne*, vol. 288, Lect. Notes. Math., Springer - Verlag, 1972.
- [K-S] M. Kashiwara and P. Schapira, *Sheaves on Manifolds*, Grund. math. Wiss. 292, Springer - Verlag, 1990.
- [La] A. Landman, *On Picard-Lefschetz Transformation for algebraic manifolds acquiring general singularities*, Trans. AMS **181** (1973), 89–126.
- [Lê1] D. T. Lê, *Une application d'un théorème d'A'Campo a l'equisingularité*, Indagat. Math. **35** (1973), 403–409.
- [Lê2] ———, *The Geometry of the Monodromy Theorem*, C. P. Ramanujam - a tribute, Tata Institute, Springer-Verlag (1978), 157–173.
- [L-M] Lê D. T. and D. Massey, *Hypersurface Singularities and the Swing*, preprint (2005).
- [Mac] R. MacPherson, *Global Questions in the Topology of Singular Spaces*, Proc. International Congress Math. 1983, Warsaw (1984), North-Holland, 213–235.
- [Mas1] D. Massey, *Lê Cycles and Hypersurface Singularities*, Lecture Notes in Mathematics, vol. 1615, Springer-Verlag, 1995.
- [Mas2] ———, *The Nexus Diagram and Integral Restrictions on the Monodromy*, to appear in J. London Math. Soc. (2004).
- [Mas3] ———, *Numerical Control over Complex Analytic Singularities*, Memoirs of the AMS, No. 778, AMS, 2003.
- [Mas4] ———, *The Sebastiani-Thom Isomorphism in the Derived Category*, Compos. Math. **125** (2001), 353–362.
- [Mi] J. Milnor, *Singular Points of Complex Hypersurfaces*, Annals of Math. Studies, no. 77, P.U.P., 1968.
- [N1] A. Némethi, *Generalized local and global Sebastiani-Thom type theorems*, Compositio Math. **80** (1991), 1–14.
- [N2] ———, *Global Sebastiani-Thom theorem for polynomial maps*, J. Math. Soc. Japan **43** (1991), 213–218.
- [O] M. Oka, *On the homotopy type of hypersurfaces defined by weighted homogeneous polynomials*, Topology **12** (1973), 19–32.
- [R] R. Randell, *On the Topology of Nonisolated Singularities*, Proc. Georgia Top. Conf., Athens, Georgia, 1977 (1979), Academic Press, New York, 445–473.

- [Sa] K. Sakamoto, *The Seifert matrices of Milnor fiberings defined by holomorphic functions*, J. Math. Soc. Japan **26** (4) (1974), 714–721.
- [Si] D. Siersma, *Variation mappings on singularities with a 1-dimensional critical locus*, Topology **30** (4) (1991), 445–469.
- [Sch] J. Schürmann, *Topology of Singular Spaces and Constructible Sheaves*, Monografie Matematyczne, vol. 63, Birkhauser, 2003.
- [S-T] M. Sebastiani and R. Thom, *Un résultat sur la monodromie*, Invent. Math. **13** (1971), 90–96.
- [T] T. Torrelli, *Intersection homology  $\mathcal{D}$ -Module and Bernstein polynomials associated with a complete intersection*, preprint (2004).

DAVID B. MASSEY, DEPT. OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MA, 02115, USA  
E-mail address: [dmassey@neu.edu](mailto:dmassey@neu.edu)