

NATURAL COMMUTING OF VANISHING CYCLES AND THE VERDIER DUAL

DAVID B. MASSEY

ABSTRACT. We prove that the shifted vanishing cycles and nearby cycles commute with Verdier dualizing up to a **natural** isomorphism, even when the coefficients are not in a field.

1. INTRODUCTION

In this short, technical, paper, we prove a result whose full statement is missing from the literature, and which may be surprising to some experts in the field. To state this result, we need to use technical notions and notations; references are [4], [2], [8], and Appendix B of [5]. We should remark immediately that the definition that we use (see below) for the vanishing cycles is the standard one, which is shifted by one from the definition in [4]

We fix a base ring, R , which is a commutative, regular, Noetherian ring, with finite Krull dimension (e.g., \mathbb{Z} , \mathbb{Q} , or \mathbb{C}). Throughout this paper, by a topological space, we will mean a locally compact space. When we write that \mathbf{A}^\bullet is complex of sheaves on a topological space, X , we mean that \mathbf{A}^\bullet is an object in $D^b(X)$, the derived category of bounded, complexes of sheaves of R -modules on X . When X is complex analytic, we may also require that \mathbf{A}^\bullet is (complex) constructible, and write $\mathbf{A}^\bullet \in D_c^b(X)$. We let $\mathcal{D} = \mathcal{D}_X$ denote the Verdier dualizing operator on $D_c^b(X)$. We will always write simply \mathcal{D} , since the relevant topological space will always be clear.

Suppose that $f : X \rightarrow \mathbb{C}$ is a complex analytic function, where X is an arbitrary complex analytic space, and suppose that we have a complex of sheaves \mathbf{A}^\bullet on X . We let ψ_f and ϕ_f denote the nearby and vanishing cycle functors, respectively. Henceforth, we shall always write these functors composed with a shift by -1 , that is, we shall write $\psi_f[-1] := \psi_f \circ [-1]$ and $\phi_f[-1] := \phi_f \circ [-1]$. In order to eliminate any possible confusion over indexing/shifting: with the definitions that we are using, if $F_{f,p}$ denotes the Milnor fiber of f at $p \in f^{-1}(0)$ inside the open ball $\mathring{B}_\epsilon(p)$ (using a local embedding into affine space), then $H^k(\psi_f[-1]\mathbf{A}^\bullet)_p \cong \mathbb{H}^{k-1}(F_{f,p}; \mathbf{A}^\bullet)$ and $H^k(\phi_f[-1]\mathbf{A}^\bullet)_p \cong \mathbb{H}^k(\mathring{B}_\epsilon(p), F_{f,p}; \mathbf{A}^\bullet)$.

The questions in which we are interested is:

Do there exist isomorphisms $\mathcal{D} \circ \psi_f[-1] \cong \psi_f[-1] \circ \mathcal{D}$ and $\mathcal{D} \circ \phi_f[-1] \cong \phi_f[-1] \circ \mathcal{D}$, even if R is not a field, and do there exist such isomorphisms which are natural?

The answer is: **yes**. In this paper, we show quickly that such natural isomorphisms exist, even if the base ring is not a field.

Is this result known and/or surprising? Some references, such as [1], [2], and our own Appendix B of [5], state that there exist non-natural isomorphisms, and require that the base ring is a field. In [8], Corollary 5.4.4, Schürmann proves the natural isomorphism exists on the stalk level, even when R is not a field.

2000 *Mathematics Subject Classification.* 32B15, 32C35, 32C18, 32B10.

Key words and phrases. vanishing cycles, nearby cycles, Verdier dual, constructible complexes.

In the l -adic algebraic context, Illusie proves in [3] that the Verdier dual and nearby cycles commute, up to natural isomorphism. M. Saito proves the analogous result in the complex analytic setting, with field coefficients, in [6] and [7]. One can obtain our full result by combining Proposition 8.4.13, Proposition 8.6.3, and Exercise VIII.15 of [4], though our proof here is completely different. In fact, our proof is similar to the discussion on duality of local Morse data following Remark 5.1.7 in [8].

Our proof is relatively simple, and consists of three main steps: proving a small lemma about pairs of closed sets which cover a space, using a convenient characterization/definition of the vanishing cycles, and using that the stratified critical values of f are locally isolated. The nearby cycle result follows as a quick corollary of the result for vanishing cycles.

We thank Jörg Schürmann for providing valuable comments and references for related results.

2. TWO LEMMAS

We shall use \simeq to denote natural isomorphisms of functors.

The following is an easy generalization of the fact that, if i is the inclusion of an open set, then $i^* \simeq i^!$ (see, for instance, Corollary 3.2.12 of [2]).

Lemma 2.1. *Suppose Z is a closed subset of X , and let $j : Z \hookrightarrow X$ denote the inclusion. Let $D_Z^b(X)$ denote the full subcategory of $D^b(X)$ of complexes \mathbf{A}^\bullet such that $Z \cap \text{supp } \mathbf{A}^\bullet$ is an open subset of $\text{supp } \mathbf{A}^\bullet$. Then, there is a natural isomorphism of functors $j^! \simeq j^*$ from $D_Z^b(X)$ to $D^b(Z)$.*

Proof. Let $\mathbf{A}^\bullet \in D_Z^b(X)$. We will show that the natural map $j^! \rightarrow j^*$ of functors from $D^b(X)$ to $D^b(Z)$ yields an isomorphism $j^! \mathbf{A}^\bullet \rightarrow j^* \mathbf{A}^\bullet$.

Let $Y := \text{supp } \mathbf{A}^\bullet$. Let $m : Y \hookrightarrow X$, $\hat{m} : Z \cap Y \hookrightarrow Z$, and $\hat{j} : Z \cap Y \hookrightarrow Y$ denote the inclusions. Then,

$$j^! \mathbf{A}^\bullet \cong j^! m_* m^* \mathbf{A}^\bullet \cong \hat{m}_* \hat{j}^! m^* \mathbf{A}^\bullet \cong \hat{m}_! \hat{j}^* m^* \mathbf{A}^\bullet \cong j^* m_! m^* \mathbf{A}^\bullet \cong j^* \mathbf{A}^\bullet,$$

where we used, in order, that $\mathbf{A}^\bullet \cong m_* m^* \mathbf{A}^\bullet$, since Y is the support of \mathbf{A}^\bullet , Proposition 2.6.7 of [4] on Cartesian squares, that $\hat{m}_* \simeq \hat{m}_!$ and $\hat{j}^! \simeq \hat{j}^*$, since \hat{m} is a closed inclusion and \hat{j} is an open inclusion, Proposition 2.6.7 of [4] again, and that $\mathbf{A}^\bullet \cong m_! m^* \mathbf{A}^\bullet$. \square

The lemma that we shall now prove certainly looks related to many propositions we have seen before, and may be known, but we cannot find a reference. The lemma tells us that, in our special case, the morphism of functors described in Proposition 3.1.9 (iii) of [4] is an isomorphism.

Lemma 2.2. *Let X be a locally compact space, and let Z_1 and Z_2 be closed subsets of X such that $X = Z_1 \cup Z_2$. Denote the inclusion maps by: $j_1 : Z_1 \hookrightarrow X$, $j_2 : Z_2 \hookrightarrow X$, $\hat{j}_1 : Z_1 \cap Z_2 \hookrightarrow Z_2$, $\hat{j}_2 : Z_1 \cap Z_2 \hookrightarrow Z_1$, and $m = j_1 \hat{j}_2 = j_2 \hat{j}_1 : Z_1 \cap Z_2 \hookrightarrow X$.*

Then, we have the following natural isomorphisms

$$(1) \quad \hat{j}_1^* \hat{j}_2^! \simeq m^* j_2! j_2^! \simeq m^* j_1^* \hat{j}_2! \hat{j}_2^! j_1^* \simeq \hat{j}_2^! j_1^*.$$

Proof. Let $i_1 : X - Z_1 \hookrightarrow X$ and $i_2 : X - Z_2 \hookrightarrow X$ denote the open inclusions. We make use of Proposition 2.6.7 of [4] on Cartesian squares repeatedly. We also use repeatedly that, if j is a closed inclusion, then $j_* \simeq j_!$ and $j^* j_* \simeq j^* j_! \simeq \text{id}$.

We find

$$m^* j_{2!} j_2^! = (j_2 \hat{j}_1)^* j_{2!} j_2^! \simeq \hat{j}_1^* \hat{j}_2^* j_{2!} j_2^! \simeq \hat{j}_1^* j_2^!,$$

which proves the first isomorphism in (1).

We also find

$$m^* j_{1*} \hat{j}_{2!} \hat{j}_2^! j_1^* = (j_1 \hat{j}_2)^* j_{1*} \hat{j}_{2!} \hat{j}_2^! j_1^* \simeq \hat{j}_2^* \hat{j}_1^* j_{1*} \hat{j}_{2!} \hat{j}_2^! j_1^* \simeq \hat{j}_2^* \hat{j}_{2!} \hat{j}_2^! j_1^* \simeq \hat{j}_2^! j_1^*,$$

which proves the last isomorphism in (1).

The middle isomorphism takes a bit more work.

As a first step, we find

$$\begin{aligned} j_{1*} \hat{j}_{2!} \hat{j}_2^! j_1^* &\simeq j_{1*} \hat{j}_2^* \hat{j}_2^! j_1^* \simeq (j_1 \hat{j}_2)^* \hat{j}_2^! j_1^* = (j_2 \hat{j}_1)^* \hat{j}_2^! j_1^* \simeq j_{2*} \hat{j}_1^* \hat{j}_2^! j_1^* \simeq \\ &j_{2*} \hat{j}_2^! j_{1*} j_1^* \simeq j_{2!} \hat{j}_2^! j_{1*} j_1^*. \end{aligned}$$

Therefore, we need to show that

$$(2) \quad m^* j_{2!} j_2^! \simeq m^* j_{2!} \hat{j}_2^! j_{1*} j_1^*.$$

Consider the natural distinguished triangle

$$i_{1!} i_1^! \rightarrow \text{id} \rightarrow j_{1*} j_1^* \xrightarrow{[1]}.$$

Applying $m^* j_{2!} j_2^!$ to this triangle, we obtain the natural distinguished triangle

$$m^* j_{2!} j_2^! i_{1!} i_1^! \rightarrow m^* j_{2!} j_2^! \rightarrow m^* j_{2!} \hat{j}_2^! j_{1*} j_1^* \xrightarrow{[1]}.$$

We claim that $m^* j_{2!} j_2^! i_{1!} i_1^! = 0$, which would yield (2) and finish the proof. We show this in two steps: we first show that $j_{2!} \hat{j}_2^! i_{1!} i_1^! \cong i_{1!} i_1^!$ and then that $m^* i_{1!} i_1^! = 0$.

Consider the natural distinguished triangle

$$j_{2!} j_2^! \rightarrow \text{id} \rightarrow i_{2*} i_2^* \xrightarrow{[1]},$$

applied to $i_{1!} i_1^!$; this yields the natural distinguished triangle

$$j_{2!} \hat{j}_2^! i_{1!} i_1^! \rightarrow i_{1!} i_1^! \rightarrow i_{2*} i_2^* i_{1!} i_1^! \xrightarrow{[1]},$$

However, $i_2^* i_{1!} = 0$, since $X = Z_1 \cup Z_2$ and, thus, we conclude that the natural map $j_{2!} \hat{j}_2^! i_{1!} i_1^! \rightarrow i_{1!} i_1^!$ is an isomorphism.

It remains for us to show that $m^* i_{1!} i_1^! = 0$, but this is trivial, because

$$m^* i_{1!} i_1^! = (j_1 \hat{j}_2)^* i_{1!} i_1^! \simeq \hat{j}_2^* j_1^* i_{1!} i_1^!,$$

and $j_1^* i_{1!} = 0$, since j_1 and i_1 are inclusions of disjoint sets. □

3. THE MAIN THEOREM

Let $f : X \rightarrow \mathbb{C}$ be complex analytic, and let $\mathbf{A}^\bullet \in D_c^b(X)$. For any real number θ , let

$$Z_\theta := f^{-1}(e^{i\theta}\{v \in \mathbb{C} \mid \operatorname{Re} v \leq 0\})$$

and let

$$L_\theta := f^{-1}(e^{i\theta}\{v \in \mathbb{C} \mid \operatorname{Re} v = 0\}).$$

Let $j_\theta : Z_\theta \hookrightarrow X$ and $p : f^{-1}(0) \hookrightarrow X$ denote the inclusions.

Following Exercise VIII.13 of [4] (but reversing the inequality, and using a different shift), we define (or characterize up to natural isomorphism) the shifted vanishing cycles of \mathbf{A}^\bullet along f to be

$$\phi_f[-1]\mathbf{A}^\bullet := p^*R\Gamma_{Z_0}(\mathbf{A}^\bullet) \simeq p^*j_{0!}j_0^!\mathbf{A}^\bullet.$$

In fact, for each θ , we define the shifted vanishing cycles of \mathbf{A}^\bullet along f at the angle θ to be

$$\phi_f^\theta[-1]\mathbf{A}^\bullet := p^*j_{\theta!}j_\theta^!\mathbf{A}^\bullet.$$

There are the well-known natural isomorphisms $\tilde{T}_f^\theta : \phi_f^0[-1] \rightarrow \phi_f^\theta[-1]$, induced by rotating \mathbb{C} counter-clockwise by an angle θ around the origin. The natural isomorphism $\tilde{T}_f^{2\pi} : \phi_f[-1] \rightarrow \phi_f[-1]$ is the usual monodromy automorphism on the vanishing cycles.

We now prove the main theorem.

Theorem 3.1. *There is a natural isomorphism of functors from $D_c^b(X)$ to $D_c^b(f^{-1}(0))$:*

$$\phi_f[-1] \circ \mathcal{D} \simeq \mathcal{D} \circ \phi_f[-1].$$

Proof. Let m denote the inclusion of $L_0 = L_\pi$ into X . Now, apply Lemma 2.2 to $X = Z_0 \cup Z_\pi$, and conclude that

$$m^*j_{0!}j_0^! \simeq \hat{j}_0^!j_\pi^*.$$

Dualizing, we obtain

$$(3) \quad \mathcal{D}(m^*j_{0!}j_0^!) \simeq \mathcal{D}(\hat{j}_0^!j_\pi^*) \simeq \hat{j}_0^!j_\pi^!\mathcal{D} \simeq m^*j_{\pi!}j_\pi^!\mathcal{D},$$

where the second isomorphism uses that \mathcal{D} “commutes” with the standard operations, and the last isomorphism results from using that $m = j_\pi\hat{j}_0$.

Let q denote the inclusion of $f^{-1}(0)$ into $L_0 = L_\pi$, so that the inclusion p equals mq . Applying q^* to (3), we obtain

$$q^*\mathcal{D}(m^*j_{0!}j_0^!) \simeq q^*m^*j_{\pi!}j_\pi^!\mathcal{D} \simeq p^*j_{\pi!}j_\pi^!\mathcal{D} = \phi_f^\pi[-1] \circ \mathcal{D} \simeq \phi_f[-1] \circ \mathcal{D},$$

where, in the last step, we used the natural isomorphism $(\tilde{T}_f^\pi)^{-1}$.

As $\mathcal{D}(q^!m^*j_{0!}j_0^!) \simeq q^*\mathcal{D}(m^*j_{0!}j_0^!)$, it remains for us to show that $q^!m^*j_{0!}j_0^!$ is naturally isomorphic to $q^*m^*j_{0!}j_0^! \simeq p^*j_{0!}j_0^! \simeq \phi_f[-1]$. This will follow from Lemma 2.1, once we show that, for all $\mathbf{A}^\bullet \in D_c^b(X)$, $f^{-1}(0) \cap \operatorname{supp}(m^*j_{0!}j_0^!\mathbf{A}^\bullet)$ is an open subset of $\operatorname{supp}(m^*j_{0!}j_0^!\mathbf{A}^\bullet)$.

Suppose that $x \in f^{-1}(0) \cap \operatorname{supp}(m^*j_{0!}j_0^!\mathbf{A}^\bullet)$. We need to show that there exists an open neighborhood \mathcal{W} of x in X such that $\mathcal{W} \cap \operatorname{supp}(m^*j_{0!}j_0^!\mathbf{A}^\bullet) \subseteq f^{-1}(0)$.

Fix a Whitney stratification of X , with respect to which \mathbf{A}^\bullet is constructible. Then, select \mathcal{W} so that all of the stratified critical points of f , inside \mathcal{W} , are contained in $f^{-1}(0)$. Suppose that there were a point $y \in \mathcal{W}$ such that $y \in f^{-1}(L_0) - f^{-1}(0)$ and the stalk cohomology of $m^*j_{0!}j_0^!\mathbf{A}^\bullet$ at y is non-zero. Then,

by definition, y would be a point in the support $\phi_{f-f(y)}[-1]\mathbf{A}^\bullet$, which, again, is contained in the stratified critical locus of f and, hence, is contained in $f^{-1}(0)$. This contradiction concludes the proof. \square

We continue to let $p : f^{-1}(0) \hookrightarrow X$ denote the closed inclusion, and now let $i : X - f^{-1}(0) \hookrightarrow X$ denote the open inclusion. Consider the two fundamental distinguished triangles related to the nearby and vanishing cycles:

$$p^*[-1] \xrightarrow{\text{comp}} \psi_f[-1] \xrightarrow{\text{can}} \phi_f[-1] \xrightarrow{[1]}$$

and

$$\phi_f[-1] \xrightarrow{\text{var}} \psi_f[-1] \rightarrow p^![1] \xrightarrow{[1]}.$$

The morphisms *comp*, *can*, and *var* are usually referred to as the *comparison map*, *canonical map*, and *variation map*. As $p^*i_! = 0 = p^!i_*$ and as $\psi_f[-1]$ depends only on the complex outside of $f^{-1}(0)$, the top triangle, applied to $i_!i^!$ and the bottom triangle applied to i_*i^* yield natural isomorphisms

$$\alpha : \psi_f[-1] \xrightarrow{\simeq} \phi_f[-1]i_!i^! \quad \text{and} \quad \beta : \phi_f[-1]i_*i^* \xrightarrow{\simeq} \psi_f[-1].$$

Corollary 3.2. *There is a natural isomorphism of functors from $D_c^b(X)$ to $D_c^b(f^{-1}(0))$:*

$$\psi_f[-1] \circ \mathcal{D} \simeq \mathcal{D} \circ \psi_f[-1].$$

Proof.

$$\psi_f[-1] \circ \mathcal{D} \simeq \phi_f[-1]i_!i^! \circ \mathcal{D} \simeq \mathcal{D} \circ \phi_f[-1]i_*i^* \simeq \mathcal{D} \circ \psi_f[-1].$$

\square

REFERENCES

- [1] Brylinski, J. L. and Monteiro Fernandes, T. *Géométrie et Analyse Microlocales*, volume 140-141 of *Astérisque*. Soc. Math. France, 1986.
- [2] Dimca, A. *Sheaves in Topology*. Universitext. Springer-Verlag, 2004.
- [3] Illusie, L. *Autour du théorème de monodromie locale*, volume 223 of *Astérisque*, pages 9–57. Soc. Math. France, 1994.
- [4] Kashiwara, M. and Schapira, P. *Sheaves on Manifolds*, volume 292 of *Grund. math. Wissen*. Springer-Verlag, 1990.
- [5] Massey, D. *Numerical Control over Complex Analytic Singularities*, volume 778 of *Memoirs of the AMS*. AMS, 2003.
- [6] Saito, M. Modules des Hodge polarisables. *Publ. RIMS*, 24 no. 6:949–995, 1988.
- [7] Saito, M. Duality for vanishing cycles functors. *Publ. RIMS*, 25 no. 6:889–921, 1989.
- [8] Schürmann, J. *Topology of Singular Spaces and Constructible Sheaves*, volume 63 of *Monografie Matematyczne*. Birkhäuser, 2004.